

JURNAL SCOPUS 2

by Artati Artati

Submission date: 24-Oct-2021 03:28PM (UTC+0700)

Submission ID: 1682306482

File name: 662-1317-1-SM.pdf (819.58K)

Word count: 7538

Character count: 35601

Open Access Article

The Effect of Variant Concentration of Kesambi (*Scheleichera Oleosa*) on Diabetic Animal Models

Rudy Hartono^{1*}, Sairah Abdul Karim², Artati^{3,4}, Arfiani Nur⁵, Ummi Zahra⁵, Aswita Amir¹

¹ Nutrition Department, Makassar Health Polytechnic, Makassar, Indonesia

² Biomedic Department, Management and Science University, Selangor Darul Ehsan, Malaysia

³ Medical Analyst Department, Makassar Health Polytechnic, Makassar, Indonesia

⁴ Chemistry Department, Faculty of Science and Math, Hasanuddin University, Makassar, Indonesia

⁵ Chemistry Department, Faculty of Science and Technology, Universitas Islam Negeri Alhoddin Makassar, Gowa, Indonesia

Abstract: Diabetes mellitus (DM) is a disease with high prevalence which continues to increase. Existing treatments, either orally or through insulin injection, has varied effectiveness for diabetics. Therefore, an alternative treatment of ⁴² is needed. Kesambi leaves have antioxidant properties, which are thought to have potential as an antidiabetic. Antioxidants are compounds that can reduce the negative ³¹ impact of free radicals and other oxidants, to inhibit oxidative damage that could worsen the severity of DM. This study aims to determine the antioxidant compound content of kesambi extract and its effects on glucose, insulin, and regeneration of pancreatic beta cells in Wistar rats induced by streptozotocin. Methods: Experimental research with modified pretest and posttest randomized controlled group design, using 60 male Wistar rats divided into 6 thin-layer ¹⁷ chromatography (TLC) groups: healthy, sick (streptozocin (STZ) induction), galvus (synthetic drug), and kesambi extract at levels of 100, 200, and 300 mg/Kg BB male Wistar rats. The study showed that kesambi (*Scheleichera oleosa*) extract had antioxidant activity of inhibitory concentration (IC) 50 of 20.29 ppm (very strong), effected the regeneration of pancreatic β cells, improved insulin levels, and decreased blood glucose levels of Wistar rats induced by STZ, dosed optimally at a concentration of 200 mg to improve regeneration of pancreatic β cells, insulin levels and decreased blood glucose levels.

Keywords: diabetes mellitus, insulin, glucose, kesambi (*Scheleichera oleosa*), β -pancreas.

不同浓度的凯桑比 (油菜花) 对糖尿病动物模型的影响

摘要: 糖尿病 (DM) 是一种患病率很高的疾病, 并且还在持续增加。现有的治疗方法, 无论是口服还是注射胰岛素, 对糖尿病患者的疗效各不相同。因此, 需要另一种 DM 治疗方法。凯桑比叶具有抗氧化特性, 被认为具有抗糖尿病潜力。抗氧化剂是可以减少自由基和其他氧化剂的负面影响 ⁴⁸ 的化合物, 以抑制可能加重 DM 严重程度的氧化损伤。本研究旨在确定凯桑比提取物的抗氧化化合物含量及其对葡萄糖、胰岛素和胰腺 β 细胞再生的影响。方法: 采用改良前测和后测随机对照设计的实验研究。使用 60 只雄性威斯塔大鼠分为 6 个薄层色谱 (薄层色谱) 组: 健康、生病 (链脲佐菌素 (STZ) 诱导)、格列齐特 (合成药物) 和凯桑比 100、200 和 300 毫克/公斤 BB 雄性威斯塔大鼠的提取物。研究证明, 凯桑比 (油菜花) 提取物具有抑制浓度 (我知道了) 50 20.29 百万分之一 (非常强) 的抗氧化活性, 影响胰腺 β 细胞的再生, 改善胰岛素水平, 降低 STZ 诱导的威斯塔大鼠

Received: April 27, 2021 / Revised: May 24, 2021 / Accepted: June 26, 2021 / Published: July 31, 2021

Notes the authors: Rudy Hartono, Nutrition Department, Makassar Health Polytechnic, Makassar, Indonesia; Sairah Abdul Karim, Biomedic Department, Management and Science University, Selangor Darul Ehsan, Malaysia; Artati, Medical Analyst Department, Makassar Health Polytechnic, Makassar, Indonesia; Chemistry Department, Faculty of Science and Math, Hasanuddin University, Makassar, Indonesia; Arfiani Nur, Ummi Zahra, Chemistry Department, Faculty of Science and Technology, Universitas Islam Negeri Alhoddin Makassar, Gowa, Indonesia; Aswita Amir, Nutrition Department, Makassar Health Polytechnic, Makassar, Indonesia

*Corresponding author. Rudy Hartono, rdharts70@gmail.com

的血糖水平, 最佳剂量为 200 毫克, 以改善胰腺 β 细胞的再生、胰岛素水平和降低血糖水平。

关键词: 糖尿病、胰岛素、葡萄糖、凯萨比 (油菜花)、 β -胰岛。

14

1. Introduction

Diabetes mellitus (DM) is a metabolic disorder occurring due to disruption in function of insulin hormone produced by the pancreas [22]. Based on 2018 data from the Indonesian Ministry of Health, Indonesia is ranked fourth of 20 countries which are estimated to have the highest number of DM cases. DM sufferers are prone to develop ROS (reactive oxygen species). The hyperglycemia state in DM triggers glucose autooxidation, resulting in ROS [1]. ROS are free radical compounds that react with other compounds. In the body, ROS tends to react with tissues and damage them. The increase of ROS in the body causes oxidative stress. Furthermore, oxidative stress contributes to islet damage and insulin resistance, which worsens the condition of DM sufferers [1].

ROS reactivity can be inhibited by antioxidants (polyphenols, flavonoid, alkaloids, and tannins) found in plants. Antioxidants can suppress beta cell apoptosis without altering the proliferation of pancreatic beta cells. Antioxidants can bind free radicals, then reduce insulin resistance [2].

Previous studies have shown a decreased level of GLUT 2 protein expression (down-regulation) in pancreatic beta cells due to the induction of streptozotocin nicotinamide, which causes toxicity to pancreatic beta cells so they are damaged and reduce insulin secretion [3]. Streptozotocin is selectively toxic to pancreatic beta cells producing insulin [4]. This compound has a glucose-like molecular shape (glucosimetic). When the compound is induced into laboratory mice, the GLUT 2 in pancreatic beta cells will be recognized as glucose. Streptozotocin will be taken to the cytosol [5]. In the cytosol, streptozotocin will undergo a redox reaction that produces ROS. The formation of ROS will cause depolarization of beta cell membranes and an increase in Ca^{2+} so the cytosol will activate various enzymes that cause lipid peroxidation, DNA fragmentation, and protein fragmentation. As a result, pancreatic beta cells become necrotic, so that insulin function and secretion decrease [6].

Treatment of DM uses hypoglycemic drugs such as injectable insulin and oral antidiabetic drugs. The administration of these drugs causes various side effects such as headaches, dizziness, nausea, and anorexia [7]. This has prompted the search for antidiabetic drugs made from natural ingredients. It is hoped that alternative natural ingredients will be more effective and have minimal side effects with the

ethnopharmaceutical and ethnobotany approach [8].

Kesambi leaves (*Schleichera oleosa*) contain secondary metabolites in the form of alkaloids, phenolics, tannins, and flavonoids. The highest levels are in phenolics and flavonoids [9]. The antioxidant activity of kesambi leaves is high, with an IC_{50} value of 16.1 ppm [10]. The high content of antioxidants in kesambi (*Schleichera oleosa*) leaves led to the development of kesambi leaf extract (*Schleichera oleosa*) into standardized herbs that have a potential anti-DM effect [11]. Previous studies have not looked at the potential of these leaves by direct indication to test animals (in vivo). The metabolic response of living beings' bodies could be proved. The novelty of this study that it is able to provide an effective antidiabetic solution from the sample so researchers will explore the effect of kesambi leaf extract (*Schleichera oleosa*) on DM using mice that will be damaged by pancreas beta cells using Streptozotocin.

2. Methods

This study used an experimental method with a modified pretest posttest randomized controlled group design, which was divided into 6 groups, each group consisting of 10 rats. The research was conducted from March to September 2020 at the Makassar Clinical Pharmacy Laboratory, Biopharmaceutical, Faculty of Pharmacy, Hasanuddin University, Makassar Faculty of Mathematics and Natural Sciences Biochemistry Laboratory, Makassar GG Laboratory, and Maros Veterinary Center.

The tools used in this study were Elisa (BIO Rad Mode 650), analytical balance (Startorius), micropipette (Ependorf), a set of minor surgical tools, a routine staining histopathologic examination tool (Haematoxylin Eosin), a maceration kit, a rotary evaporator (Ruchi), magnetic stirrer, UV-Vis spectrophotometer (Hewlett Packard), animal scale, centrifuge, syringe, oral needle, binocular microscope, camera and mouse cage.

The materials used in this study were kesambi leaves taken from Barru Regency, South Sulawesi. White wistar male rats were obtained from the Biopharmaceutical Laboratory of the Faculty of Pharmacy, UNHAS. Other materials were Folin-Ciocalteu reagent, pH 4.5 citrate buffer, distilled water, nqua pro injection (Otsuka), alcohol (E. Merck), buffered formaldehyde, xylol, paraffin, diethyl ether (E. Merck), Na. CMC, STZ, ALX-350-010 from

ALEXIS Corporation, animal feed (HG II-B), histopathologic material, Haematoxylin Eosin staining and immunohistochemical staining, Bovin Serum Albumin (BSA), KCl, KH_2PO_4 , NaCl, $\text{Na}_2\text{HPO}_4 \cdot \text{H}_2\text{O}$, NaOH, Tween, NaNO_3 (E. Merck), Antibodies, and mouse insulin.

The research sample was obtained from the population by simple random sampling with inclusion criteria (white male rats, strain wistar, age 1.5-2 months, body weight 150-200 grams. During adaptation 14 days before treatment, all rats were not sick; their activity and behavior were normal). Sixty rats were used in the research.

2.1. Kesambi Leaf Extraction

The leaves of Kesambi (*Schleichera oleosa*) were cleaned and dried. A total of 5.90 kg of sample was extracted with n-hexane solvent. Then the phytochemical test was carried out using FeCl_3 , bromine water, and indophenol. Quantitative analysis of polyphenols [29] performed using the Folin-Ciocalteu method at a wavelength of 666 nm using a UV-Vis spectrophotometer.

2.2. Antioxidant Activity Test

Two ml of sample solution of various concentrations were added with 2 ml of DPPH 0.4 mM and left at room temperature for 30 minutes. Then it was measured on a visible spectrophotometer at a wavelength of 515 nm. The IC_{50} value is determined by probit between the log data of the concentration and the probit percentage of free radical binding.

2.3. Application to Diabetic Animal Models

Kesambi extract suspension was made: 100 mg/kg mass of rats and 300 mg/kg BW of rats, respectively. Sixty rats that met the inclusion criteria were taken randomly on day 0 and then put into 6 groups. Each group consisted of 10 rats and was kept in separate pens, given standard food and drink ad libitum.

Furthermore, each group was given the following treatment respectively. All groups consisting of ten rats were fasted first and observed. *Group I* was healthy control rats; *Group II* the rats were induced STZ 60 mg/kg single dose rats as a pain control; *Group III* the rats were induced STZ 60 mg/kg single dose rats by i.p. as a positive comparison. Starting from the seventh day, the rats were given 0.9 mg/200g gnlvus orally every day for 21 days. In *Groups IV-VI*, the rats were induced STZ 60 mg/kg [30] single dose rats by i.p. Starting from the seventh day 100 mg/kg, 200 mg/kg, and 300 mg/kg of rat kesambi extract were given orally every day for 21 days.

Histopathologic examination of pancreatic β cells, insulin levels and blood glucose levels were performed five times; once before STZ induction, and four times after being induced by STZ, on day 1, 7, 14, 21, and

28. Pancreatic tissue was taken and a blood smear was then performed for histopathologic examination by staining using hematoxylin eosin to [34]nitatively examine the number of pancreatic β cells. Insulin levels were measured using Enzyme Linked Immuno Sorbent Assay (ELISA). Blood glucose levels were measured using a glucometer.

2.4. Data Analysis

Data analysis was performed using the ANOVA test to compare the effect of [5]eral treatment interventions on blood glucose levels, pancreatic β cell counts, and insulin levels in 6 groups of male Wistar rats.

3. Results and Discussion

The extract was tested by adding a specific reagent for alkaloids, flavonoids, steroid [47]annin and saponin. Results of phytochemistry assay are shown in Table 1.

Table 1. Phytochemistry assay

Phytochemistry assay	Result
Steroid	+
Alkaloid	+
Flavonoid	+
Saponin	+
Tannin	+

The results showed that the extract contained a secondary metabolite that has potential as antioxidant agent. The results of this study were consistent with several previous studies [4], [10], [12]. An antioxidant assay was then done to determine antioxidant activity.

3.1. Antioxidant Activity [66]

Testing to determine total phenolic content for antioxidant activity w[3] performed using the Folin-Ciocalteu method. The antioxidant activity was carried out using the DPPH method. The principle of this method is the donation of hydrogen atoms ($\text{H}^{\bullet}$) from the substance tested to the DPPH radical to become a non-radical compound of dibenzyl picryl hydrazine, which is indicated by a color change. The color change that occurs is a color change from purple to yellow; the intensity of the DPPH color change is directly proportional to the antioxidant activity to reduce these free radicals. The IC_{50} results showed antioxidant activity of 20.29 ppm better than its bark stem [13]. From these results, it can be seen that ke[49]mbi leaves have very strong antioxidant activity. The IC_{50} value of less than 50 ppm has a very strong antioxidant activity. The active compound contained in the extract is a group of antioxidant compounds.

3.2. Acclimatization Test

Male rats were selected as the test animals because their biological conditions are more stable than those of female rats. Moreover, they are not influenced by the oestrous cycle and they have a faster drug metabolism

rate. The rats were acclimatized to the laboratory for 14 days, and they were given pellet feed and drinking water ad libitum.

In the acclimatization process, the rats were weighed to determine the treatment dose to be administered to them. The rats used in the study were healthy, with signs of non-standing hair, white fur, clear eyes, normal behavior, and their body weight increased during the intervention period according to the research method.

Kesambi extract is an ingredient given to the subjects of treatment given in ad libitum. To minimize the variability between the test animals, the test animals had a uniform body weight of 200–300 g and they were between three and four months of age. The selection of rats as test animals was due to their high availability and sensitivity to represent humans in determining blood glucose levels. Meanwhile, rats have metabolic and digestive systems that are similar to those of humans.

The mechanism of STZ in pancreatic cells can affect glucose causing hyperglycemia as it enters the pancreatic β cells through the glucose transporter (GLUT-2), thereby causing DNA alkylation. Alkylation or the entry of the methyl group from STZ into the DNA molecule will cause damage to the DNA fragmentation. Subsequently, the DNA damage will trigger the production of synthetic poly (ADP-ribose) enzymes, which are enzymes needed to repair the DNA damage. These enzymes require nicotinamide adenine dinucleotide (NAD) as a substrate so the NAD⁺ content in cells decreases, which causes a decrease in the amount of ATP. Thus, insulin synthesis and secretion are inhibited, thereby causing hyperglycemia [9].

According to the World Health Organization, the normal blood glucose level before meals is approximately 70–130 mg/dL. After meals, the blood glucose level naturally increases; however, it should still be less than 180 mg/dL. Additionally, the blood glucose level during fasting is less than 100 mg/dL. A synthetic drug in the form of galvus was administered to the rats in the third group in order to increase their insulin secretion and ensure a normal blood glucose level. Furthermore, kesambi extract with various doses was administered to the rats in the fourth, fifth, and sixth groups [14].

3.3. ANOVA Test Using SPSS for Windows Software

3.3.1. Glucose Levels

In Table 2, there was a difference in the average blood glucose level with a P value < 0.05. This indicates that the second group experienced an increase in blood glucose level. In this intervention, H_0 was accepted and H_a was rejected so that there is induction

effect of STZ, the administration of kesambi extract 100, 200, and 300 mg on blood glucose levels.

For the second group induced by STZ and the third group, there was no difference in the average blood glucose levels. This indicates that with the induction of STZ, the blood glucose levels of mice in the third group increased to normal values, indicating that the addition of galvus did not lead to a decrease in the blood glucose levels of the second group.

The results of the average glucose level in the third (with galvus), first, fourth, and fifth groups showed that each group had a P value < 0.05. This shows that the third group produced blood glucose levels that were close to those of normal mice (NaCMC administration) so there is an effect of intervention giving galvus towards normal blood glucose levels. For the sixth group, which was given 300 mg of kesambi extract, there was no difference in the average blood glucose level with a P value > 0.05.

From the results of the average glucose levels of the third, fourth, and fifth groups, H_0 was accepted and H_a was rejected, so that the administration of 100, 200, and 300 mg of kesambi extract with NaCMC did not have any effect. Meanwhile, there was a difference in the average blood glucose levels of the sixth group and the first group (given NaCMC) after the ANOVA test. In this case, the effect of the administration of 300 mg of kesambi extract with NaCMC was observed.

Table 2. Post hoc test for glucose levels

Intermediate glucose levels	(I) Group	(J) Group	Mean Difference (I-J)	Std. Error	Sig.
Group II STZ with several KLP	STZ	NaCMC	81.87200	11.35778	.000
		Galvus	58.17600	11.35778	.015
		100 mg	66.94800	11.35778	.000
		200 mg	97.04000	11.35778	.000
		300 mg	90.94800	11.35778	.000
Group III Galvus with several KLP	Galvus	NaCMC	61.50200	11.35778	.000
		100 mg	47.77600	11.35778	.000
		200 mg	77.87000	11.35778	.000
		300 mg	58.28400	11.35778	.010
Group IV Kesambi extract 100 mg with several group	100 mg	NaCMC	54.50200	11.35778	.024
		200 mg	100.84000	11.35778	.000
		300 mg	101.00200	11.35778	.000
Group V Kesambi extract 200 mg with several group	200 mg	NaCMC	117.88800	11.35778	.000
		300 mg	42.90800	11.35778	.001

3.3.2. Insulin Levels

The results of the average insulin levels showed a difference between the average insulin levels of second group and the first and fifth groups after the ANOVA test, with a P value < 0.05. This indicates that the second group had abnormal insulin levels because STZ acts as a destroyer of the pancreatic β cell structure, which affects insulin production. However, the addition

of kesambi extract as much as 200 mg causes the production of the hormone insulin at normal levels. In this intervention, H_a was accepted and H_0 was rejected so that the effects of STZ induction and the addition of 200 mg of kesambi extract on the insulin levels were observed. In contrast, the third, fourth, and sixth groups had P values > 0.05. This shows that the third group did not have normal insulin levels. In this intervention, the addition of galvus and 100 and 300 mg of kesambi extract had no effect on the normal insulin level, as presented in Table 3.

The results of the average insulin levels between KLP III (given galvus) and KLP I (given NaCMC) after the ANOVA test showed a difference in the average insulin levels with a P value < 0.05. This shows that the third group had normal insulin production, and the effect of galvus intervention on normal insulin levels was observed. Meanwhile, in the fourth, fifth, and sixth groups with P values > 0.05, H_0 was accepted and H_a was rejected. There was no effect of the intervention between galvus and kesambi extracts of 100, 200, and 300 mg. The same thing was also found in the first, fourth, fifth, and sixth groups, as shown in Table 3.

Table 3 Post hoc test for insulin in rat

Insulin Level	(1) Group	(2) Group	Mean Difference (2-1)	Std. Error	Sig.
Group II STZ with control Group	STZ	NaCMC	42.34000	11.33482	.001
		Galvus	937000	11.33390	.348
		100 mg	14.30000	11.33382	.302
		200 mg	33.90000	11.33382	.004
		300 mg	22.70000	11.33382	.098
Group III Galvus with control Group	Galvus	NaCMC	15.70000	11.33392	.384
		100 mg	11.80000	11.33382	.179
		200 mg	24.30000	11.33382	.004
		300 mg	14.30000	11.33382	.122
KLP IV Kesambi extract 100 mg with control group	100 mg	NaCMC	14.30000	11.33382	.302
		200 mg	32.00000	11.33382	.004
		300 mg	28.10000	11.33382	.004
KLP V Kesambi extract 200 mg with group NaCMC	200 mg	NaCMC	22.30000	11.33382	.001
		300 mg	41.90000	11.33382	.001

17

3.3.3. Number of β Pancreatic Cells

The mean number of pancreatic beta cells in the first to the sixth group with all treatments after the ANOVA test showed no difference in the average number of pancreatic beta cells with a P-value > 0.05. In this intervention, H_0 is accepted, H_a is rejected so that there is no effect of the intervention between each treatment. This can be seen in Table 4.

Table 4 Mann-Whitney test for number of β -pancreatic cells

	Group	N	Mean Rank	Sum of Ranks	Asymp. Sig. (2-tailed)
Number of β -Pancreatic Cells on group I with NaCMC	KLP II	5	8.00	40.00	.008
	STZ	5	33.00	165.00	
	KLP III	5	6.00	30.00	.008
	Galvus	5	7.00	35.00	
	KLP IV	5	6.00	30.00	.008
	Kesambi extract 100 mg	5	7.00	35.00	
Number of β -Pancreatic Cells on group II with STZ	KLP V	5	7.40	37.00	.047
	Galvus	5	7.80	39.00	
	KLP VI	5	6.00	30.00	.008
	NaCMC	5	7.00	35.00	
	Kesambi extract 200 mg	5	7.00	35.00	
	Kesambi extract 300 mg	5	7.00	35.00	
Number of β -Pancreatic Cells on group III with Galvus	KLP III	5	4.50	22.50	.002
	STZ	5	8.20	41.00	
	KLP IV	5	5.60	28.00	.002
	STZ	5	8.50	42.50	
	KLP V	5	5.80	29.00	.008
	STZ	5	7.20	36.00	
Number of β -Pancreatic Cells on group IV with Kesambi extract	KLP VI	5	4.50	22.50	.002
	STZ	5	8.50	42.50	
	Kesambi extract 100 mg	5	5.80	29.00	.008
	STZ	5	7.20	36.00	
	KLP V	5	5.80	29.00	.008
	Kesambi extract 200 mg	5	7.20	36.00	
Number of β -Pancreatic Cells on group V with Kesambi extract	KLP VI	5	5.80	29.00	.008
	STZ	5	7.20	36.00	
	Kesambi extract 100 mg	5	5.80	29.00	.008
	STZ	5	7.20	36.00	
	Kesambi extract 200 mg	5	5.80	29.00	.008
	Kesambi extract 300 mg	5	5.80	29.00	.008

3.4. In-Vivo Kesambi Extract Activity in Wistar Rats on Blood Glucose Levels

There is a positive correlation between the treatment dose and the average blood glucose level of the rats given the extract of kesambi. This explained that the mean blood glucose levels continued to decrease with the administration of kesambi extract. For determining the effect of kesambi extract on decreasing blood glucose levels and which groups had significant differences, an analysis was performed using an unpaired t test.

The mean blood glucose levels of Wistar rats in the second group were significantly different from the first and fifth groups, with p-values < 0.05. They were not significantly different between the second, third, fourth groups, with p-values > 0.05. This shows that the administration of kesambi extract at a concentration of 200 mg affects reducing blood glucose levels in rats induced by STZ (H_0 is rejected, and H_a is accepted).

In the second group, after being induced by STZ starting on days 7, 14, 21, and 28, there was an increase in blood glucose levels. In the third group, after being given galvus synthetic drug on days 7, 14, 21, and 28, the sugar levels began to decrease compared to the second group, but glucose levels remained high. In the fourth, fifth, and sixth groups, each rat was given

kesambi extract with a concentration of 100, 200, and 300 mg/kg of rats on days 7, 14, 21, and 28. There was a decrease in blood glucose levels. In the fifth group, there was a significant decrease in blood glucose levels towards normal glucose levels close to glucose levels of the first group. At a concentration of 200 mg, the extract worked optimally to reduce blood glucose levels in Wistar rats that had experienced hyperglycemic due to STZ induction. This can be because kesambi leaf extract and galvus have suppressed the increase in blood glucose levels by activating pancreatic β cells for insulin production (Fig. 1).




Fig. 1 Response to glucose levels

Insulin has four effects that can lower blood glucose levels and increase carbohydrate storage. Among others, insulin makes it easier for glucose to enter most cells. Glucose molecules do not easily penetrate cell membranes in the absence of insulin. Thus, most tissues rely heavily on insulin to absorb glucose from the blood and use it. Insulin increases the facilitated diffusion (mediated by carriers) of glucose into glucose-dependent cells through the transporter recruitment phenomenon [15].

Antioxidants are thought to reduce blood glucose levels by inhibiting glucose absorption from the gastrointestinal lumen [10], increasing glucose utilization in peripheral tissues, thus acting directly on pancreatic β cells by triggering the activation of the cAMP signal cascade (cyclic Adenosine Monophosphate) in strengthening insulin secretion which is sensitive to glucose.

Antioxidants protect pancreatic β cells from the toxic effects of free radicals produced under chronic hyperglycemia conditions. The administration of antioxidants can increase the mass of pancreatic β cells and maintain the insulin content in them. In cells with insulin receptors (muscle cells, adipose cells, and liver cells), the binding of free radicals will increase insulin signaling in intracellular GLUT4 translocation to the cell membrane; then, they will be able to take glucose from the blood [6]. In general, reducing oxidative stress can reduce insulin resistance and inhibit β -pancreatic cell damage [13].

Antioxidants can prevent complications or progression of diabetes mellitus by cleaning excessive free radicals, breaking the chain of free radical reactions [16], binding metal ions (chelating), and blocking polyol pathways by inhibiting the enzyme aldolase reductase [16]. Antioxidants also have an inhibitory effect on the alpha glucosidase enzyme through hydroxylation and substitution in the β ring [17]. The principle of this inhibition is similar to the acarbose which has been used as a drug for the treatment of diabetes mellitus. This is done by producing a procrastination in the hydrolysis of carbohydrates and disaccharides, absorption of glucose, and inhibiting the metabolism of sucrose to glucose and fructose [18].

The antihyperglycemic effect is thought to be the effect of the secondary metabolite content of the antioxidant extract of kesambi. Antioxidants have a hypoglycemic effect with several mechanisms: inhibiting the glucose absorption, increasing glucose tolerance, stimulating insulin release or acting like insulin, increasing glucose uptake by peripheral tissues and regulating enzymes that play a role in carbohydrate metabolism [19].

3.5. The Activity of Kesambi Extract in Vivo in Wistar Strain Rats on Insulin Levels

The relationship between the treatment dose and the average insulin levels in the blood of rats will be demonstrated after giving them kesambi extract. It explains that the mean insulin levels decreased with the administration of kesambi extract. To determine the effect of kesambi extract on decreasing insulin levels and which groups had significant differences, an analysis was performed using the unpaired t test. The results of the mean blood glucose levels of Wistar rats at the second group were significantly different from the first and fifth group with p values <0.05 . However, there was no significant difference between the second, third, and fourth, with p values >0.05 . This shows that giving the kesambi extract at a concentration of 200 mg has an effect on decreasing the blood insulin levels in STZ-induced rats (H_0 is rejected and H_a is accepted).

At the second group, after being induced with STZ starting on days 7, 14, 21, and 28, there was an increase in the insulin levels. In the third group, after being given the synthetic drug galvus on days 7, 14, 21 and 28, the insulin levels started to decrease as compared to the second group, but the levels remained high. In the fourth, fifth, and sixth groups, the extract was given kesambi with a concentration of 100, 200 and 300 mg/kg of rats on days 7, 14, 21, and 28, in which there was a decrease in blood insulin levels. In the fifth group, there was a significant decrease in blood insulin toward normal levels approaching the first group's insulin levels. At a concentration of 200 mg, kesambi extract worked optimally to reduce blood insulin levels

in Wistar-strain rats that had experienced hyperglycemia due to STZ induction [20].

Insulin levels increased after STZ induction. This is due to the destruction of pancreatic β cells. In pancreatic β cells, the STZ reaction will cause the oxidation of the peptide glutathione in the cells and form dialuric acid, which can produce free radicals in the form of hydrogen peroxide (H_2O_2), superoxide anion (O_2^-), nitric oxide (NO), thus damaging the DNA chain of pancreatic β cells. Damage to the DNA of pancreatic β cells causes necrosis and results in problematic insulin production and secretion [10].




Fig. 2 Insulin level response

Treatment with kesambi extract is thought to play a role in restoring impaired glucose homeostasis by lowering blood glucose levels and tends to increase plasma insulin levels. The active compound extract of kesambi that plays a role in blood glucose homeostasis is unknown, but it is suspected that the acetone water-soluble compound is an unknown antioxidant group. The active compounds of this antioxidant class act as antioxidants that can prevent and reduce free radicals by reacting directly to these free radicals [21].

3.6. In Vivo Activity of Kesambi Extract in Wistar-Strain Rats on the Number of Pancreatic β Cells

The relationship between treatment dose and the number of pancreatic β cells in rats given the extract of kesambi shows that the number of pancreatic β cells affects the administration of kesambi extract. To determine the effect of kesambi extract on the number of pancreatic β cells and which groups had significant differences, an analysis was performed using an unpaired t-test.

The results of the mean number of pancreatic β cells in Wistar strain mice at the second group were significantly different from the first and fifth groups, with p values <0.05 , but between groups 2, 3, and 4 the numbers were not significantly different, with p values >0.05 . This shows that giving kesambi extract at a concentration of 200 mg affects the number of pancreatic β cells in mice induced with STZ. (H_0 is rejected and H_a is accepted.)

STZ causes damage to pancreatic β cells. One of the

STZ mechanisms causes hyperglycemia related to the formation of free radicals, including NO, O_2^- , and H_2O_2 , which can cause cell DNA fragmentation due to STZ cytotoxics. STZ has been shown to increase reactive oxygen species (ROS) that contribute to DNA fragmentation and trigger changes in cells. STZ selectively accumulates in pancreatic β cells through the low-affinity GLUT2 glucose transporter on the plasma membrane. The entry of the methyl (alkylation) group in STZ into the DNA molecule will cause damage to the DNA fragment. The DNA damage will activate cellular poly adenosine diphosphate (ADP) - nicotinamide adenine dinucleotide (NAD $^+$), further reducing adenosine triphosphate (ATP) [10]. This increase in ATP dephosphorylation will provide a substrate for xanthine oxidase and will speed up the reaction to give the superoxide anion as the end product. Activation of hydrogen peroxide and other hydroxyl radicals will result from the formation of this superoxide anion.

The infected control group (the second group) showed the occurrence of necrosis and degeneration as indicated by the decreased number of pancreatic β cells. This is because the second group did not receive an extract of kesambi leaves (*Scheuchzeria oleosa*). Whereas in the third, fourth, fifth, and sixth group, the number of pancreatic β cells increased on days 7, 21 and 28. It is suspected that the increased number of pancreatic β cells occurred due to the administration of extracts of kesambi (*Scheuchzeria oleosa*), which contains antioxidants. In the fifth group, there was a significant increase in value, with numbers approaching those of the first group, namely the healthy group (Fig. 3).




Fig. 3 Response of pancreatic β cells

Langerhans Island is formed from ductal precursor cells and can reduce oxidative stress and help restore the function of pancreatic β cells. Antioxidants help regenerate pancreatic β cells, reduce oxidative stress, and show the incidence of diabetes through reducing fat levels and increasing levels of Superoxide Dismutase (SOD), catalase and glutathione peroxidase. Antioxidants can play a role in capturing free radicals or function as natural antioxidants [10]. This antioxidant activity allows antioxidants to capture or

neutralize free radicals (such as ROS or RNS) associated with phenolic OH groups so that they can improve the state of damaged tissue; in other words, the inflammatory process can be inhibited [10].

Antioxidants that play a role in reducing blood glucose levels will increase pancreatic beta cell count, insulin secretion and sensitivity levels, and the production of alpha-glucose inhibitors. The intended antioxidant compounds that play a role in lowering blood glucose levels are flavonoid polyphenols compounds, saponins, and tannins, as reactive oxygen species (ROS). Flavonoids can capture free radicals directly through the donation of hydrogen atoms that are on the OH group and conjugate the double bond. Free radicals are made inactive. Polyphenols act as antioxidants because they have hydroxyl groups that can donate hydrogen atoms to free radical compounds and stabilize reactive oxygen compounds (ROS), and have hydroxyl ketone groups that can act as metal chelates [10].

Tannins and saponins function as inhibitors of the activity of the alpha-glucosidase enzyme, which is an enzyme in digestion that converts carbohydrates into glucose, before glucose absorption is reduced [23].

4. Conclusion

A kesambi extract (*Schleichera oleosa*) has a very strong antioxidant activity. The administration of a kesambi extract (*Schleichera oleosa*) affected the regeneration of pancreatic cells, improved insulin levels, and decreased blood glucose levels of STZ-induced Wistar rats. There is an effect of administering a graded dose of a kesambi extract (*Schleichera oleosa*) to pancreatic cell regeneration, i.e., increasing insulin levels and lowering blood glucose levels of STZ-induced Wistar rats; it works optimally on kesambi leaf extract with a concentration of 200 mg. Further investigations should be carried out on kesambi extracts; therefore, an isolation process is needed. By knowing the content of pure compounds, it is possible to know the interactions as antidiabetic drugs.

Acknowledgment

The authors thank the Ministry of Health of the Republic of Indonesia through overseas collaborative research for the financial support in this research; the Maros Veterinary Center for facilitating several tests in this research.

References

[1] AYODEJI D. O., MARYANN E. C., SAMUEL C., ALUMBUGU T. P., AUDU D. C., and ADEBAYO O. O. Dependence of ageing on reactive oxygen species production and the cushioning effect of antioxidants (review). *Pharmacologyonline*, 2019, 1: 397-402. https://pharmacologyonline.silae.it/files/archives/2019/vol1/PhOL_2019_1_A040_Durojaye.pdf

[2] BAJAJ S., & KHAN A. Antioxidants and diabetes. *Indian Journal of Endocrinology and Metabolism*, 2012, 16(Supplement 2): 267-271. <https://doi.org/10.4103/2230-8210.104057>

[3] SHOLIKAH T. A., WULANDARI S., ARIESTA I., HAKIM M. A. R., and HAFIZHAN M. The hypoglycemic effects of tapak liman (*Elephantopus scaber* L.) plant extract on albino rat (*Rattus norvegicus*) models of diabetes mellitus. *Indonesian Journal of Medicine and Health*, 2020, 11(2): 172-179. <https://doi.org/10.20885/IJKI.Vol11.Iss2.art10>

[4] SETIAMEANG R., NURANI W., IBRAHIM A. M., and SILABAN S. Analysis of secondary metabolic compounds from leaves extract kesambi (*Schleichera oleosa*) and antioxidant activity test. *Jurnal Pendidikan Kimia*, 2016, 8(3): 164-168. <https://doi.org/10.24114/jpkim.v8i3.4479>

[5] MITTLER R. ROS Are Good. *Trends in Plant Science*, 2017, 22(1): 11-19. <https://doi.org/10.1016/j.tplants.2016.08.002>

[6] IGHOJARO O. M., ADEHUN A. M., and AKINLOYE O. A. Alloxan-induced diabetes, a common model for evaluating the glycemic-control potential of therapeutic compounds and plants extracts in experimental studies. *Medicina*, 2017, 53(6): 365-374. <https://doi.org/10.1016/j.medci.2018.02.001>

[7] CHAUDHURY A., DUVOOR C., REDDY DENDI V. S., KRALEFI S., CHADA A., RAWJIA R., MARICI A., SHEKHAWAT N. S., MONTALES M. T., KURIAKOS K., SASAPU A., BEEDE A., PATIL N., MUSHAM C. K., LOHANI G. P., and MIRZA W. Clinical Review of Antidiabetic Drugs: Implications for Type 2 Diabetes Mellitus Management. *Families in Endocrinology*, 2017, 8: 6. <https://doi.org/10.3389/fendo.2017.00006>

[8] KOOTI W., FAROKHIPOUR M., ASADZADEH Z., ASHTARY-LARKY D., and ASADI-SAMANI M. The role of medicinal plants in the treatment of diabetes: a systematic review. *Electronic Physician*, 2016, 8(1): 1832-1842. <https://doi.org/10.19082/1832>

[9] PANCHE A. N., DIWAN A. D., and CHANDRA S. R. Flavonoids: An overview. *Journal of Nutritional Science*, 2016, 5: e47. <https://doi.org/10.1017/jns.2016.41>

[10] HOLIL K., & GRIANA T. P. Analisis Fitokimia dan Aktivitas Antioksidan Ekstrak Daun Kesambi (*Schleichera oleosa*) Metode DPPH. *Journal of Islamic Pharmacy*, 2020, 5(1): 28-32. <https://doi.org/10.18860/jip.v5i1.9387>

[11] GOSWAMI S., & SINGH R. P. Antidiabetic Potential and HPTLC Fingerprinting of *Schleichera oleosa* (Lour) Oken. *Pharmacognosy Journal*, 2019, 11(3): 469-474. <https://doi.org/10.5530/pj.2019.11.74>

[12] HANIFAH I. The Influence of Kesambi Leaf Extract (*Schleichera oleosa*) on Proliferation Granulosa Cell Goat Ovarium (*Capra aegagris* HIRCL.S) in Vitro. *Jurnal Biota*, 2020, 6(1): 19-25. <https://doi.org/10.19109/biota.v6i1.4372>

[13] DEWI Y. K. Uji Aktivitas Antioksidan Dari Ekstrak Kulit Batang Kesambi [*Schleichera Oleosa* (Lour) Oken] Menggunakan Metode Ekstraksi Bertingkat. *SPIN: Jurnal Kimia & Pendidikan Kimia*, 2021, 3(1): 22-31. <https://doi.org/10.20414/spin.v3i1.3020>

[14] YANG WANG T., LI Q., and SHUN BI K. Bioactive Flavonoids in medicinal plants: Structure, activity and biological fate. *Asian Journal of Pharmaceutical Sciences*, 2018, 13(1): 12-23. <https://doi.org/10.1016/j.ajps.2017.08.004>

- [15] CARITÀ A. C., FONSECA-SANTOS B., SHULTZ J. D., MICHNIAK-KOHN B., CHORILLI M., and LEONARDI G. R. Vitamin C: One compound, several uses. *Advances for delivery, efficiency and stability. Nanomedicine: Nanotechnology, Biology and Medicine*, 2020, 24: 102117. <https://doi.org/10.1016/j.nano.2019.102117>
- [16] IFEANYI O. E. A Review on Free Radicals and Antioxidant. *International Journal of Current Research in Medical Sciences*, 2018, 4(2): 123-133. <http://dx.doi.org/10.22192/ijcrms.2018.04.02.019>
- [17] ALKADI H. A Review on Free Radicals and Antioxidants. *Infectious Disorders - Drug Targets*, 2018, 20(1): 16-26. <http://dx.doi.org/10.2174/1871526518666180628124323>
- [18] MCIVER L. A., & TRIPP J. *Continuing Education Activity*. StatPearls Publishing, National Center for Biotechnology Information, 2021. <https://www.ncbi.nlm.nih.gov/books/NBK493214>
- [19] VIEIRA R., SOUJO S. H., SÁNCHEZ-LÓPEZ F., MACHADO A. L., SEVERINO P., JOSE S., SANTINI A., FORTUNA A., GARCIA M. L., SILVA A. M., and SOUJO B. B. Sugar-lowering drugs for type 2 diabetes mellitus and metabolic syndrome — review of classical and new compounds. Part-I. *Pharmaceutics*, 2019, 12(4): 152. <https://doi.org/10.3390/ph12040152>
- [20] GOSWAMI S., & SINGH R. P. Ayurvedic, Phytochemical and Pharmacological Review of *Schleichera Olsea* (Lour.) Oken: a Traditional Plant With Enormous Biological Activity. *World Journal of Pharmaceutical Research*, 2017, 6(10): 295-309. <https://doi.org/10.20959/wjpr201710-9370>
- [21] JOSE S., & SINHA M. P. A Comparative Analysis of Antioxidant Capacity of Aqueous and Methanolic Leaf Extracts of *Scoparia Dulcis* and *Schleichera Olsea*. *Balneo Research Journal*, 2018, 9(3): 221-227. <https://doi.org/10.12680/balneo.2018.185>
- [22] SHINAIN S. A. H., NABAT Z. M., SALIH M. A., and AL KHAFAJI B. J. Diagnosing Diabetes with Extended Binary Cuckoo Search and k-Nearest Neighbor Classifier. *Journal of Southwest Jiaotong University*, 2020, 55(2). <https://doi.org/10.35741/jssn.0258-2724.55.2.36>
- [23] ŞÖHRETOĞLU D., & SARI S. Flavonoids as α -glucosidase inhibitors: mechanistic approaches merged with enzyme kinetics and molecular modelling. *Phytochemistry Reviews*, 2020, 19(5): 1081-1092. <https://doi.org/10.1007/s11101-019-09610-6>
- 参考文献:
- [1] AYODEJI D. O., MARYANN E. C., SAMUEL C., ALUMBUGU T. P., AUDU D. C. 和 ADEBAYO O. O. 老化对活性氧产生的依赖性抗氧化的缓冲作用 (评论). *药理学在线*, 2019, 1: 397-402. https://pharmacologyonline.scit.az/files/archives/2019/vol1/PhOL_2019_1_A040_Durojaye.pdf
- [2] BAJAJ S., & KHAN A. 抗氧化剂和糖尿病. *印度内分泌与代谢杂志*, 2012, 16 (增刊 2): 267-271. <https://doi.org/10.4103/2230-8210.104057>
- [3] SHCHIKAH T. A., WILLANDARI S., ARIESTA I., HAKIM M. A. R. 和 HAFIZHAN M. 塔纳里曼(象牙树) 植物提取物对自化大鼠 (高血糖) 糖尿病模型的降血糖作用. *印度西医学与健康杂志*, 2020, 11(2): 172-179. <https://doi.org/10.20885/JKKI.Vol11.Iss2.art10>
- [4] SITIMFANGI R., NURAFENI W., IBRAHIM A. M. 和 SILABAN S. 叶提取物(萨比油菜花) 次生代谢物的分析和抗氧化活性测试. *期刊彭迪次基米亚*, 2016, 8(3): 164-168. <https://doi.org/10.24114/jpkim.v8i3.4479>
- [5] MITTLER R. 活性氧很好. *植物科学进展*, 2017, 22(1): 11-19. <https://doi.org/10.1016/j.plants.2016.08.002>
- [6] IGHODARO O. M., ADEOSUN A. M. 和 AKINLOYE O. A. 白鼠糖原诱导的糖尿病, 这是一种在实验研究中评估治疗性化合物和植物提取物的血糖控制能力的常用模型. *医学*, 2017, 53(6): 365-374. <https://doi.org/10.1016/j.medic.2018.02.001>
- [7] CHAUDHURY A., DUVOOR C., REDDY DENDI V. S., KRALETI S., CHADA A., RAVILLA R., MARCO A., SHEKHAWAT N. S., MONTALES M. T., KURIAKOSE K., SASAPU A., BEEBE A., PATIL N., MUSHAM C. K., LOHANI G. P., 和 MIRZA W. 抗糖尿病药物的临床评价: 对 2 型糖尿病管理的影响. *内分泌学前沿*, 2017, 8: 6. <https://doi.org/10.3389/fendo.2017.00006>
- [8] KOOTI W., FAROKHIPOUR M., ASADZADEH Z., ASHTARY-LARKY D. 和 ASADI-SAMANI M. 药用植物在糖尿病治疗中的作用: 系统评价. *电子医药*, 2016, 8(1): 1832-1842. <https://doi.org/10.19082/1832>
- [9] PANCHE A. N., DIWAN A. D. 和 CHANDRA S. R. 天然化合物: 概述. *营养科学杂志*, 2016, 3: e47. <https://doi.org/10.1017/jns.2016.41>
- [10] HOLIL K., & GRIANA T. P. 蔬菜叶提取物 (油菜花) DPPH 的植物化学分析和抗氧化活性. *方法. 伊斯兰药理学杂志*, 2020, 5(1): 28-32. <https://doi.org/10.18860/jip.v5i1.9387>
- [11] GOSWAMI S., & SINGH R. P. 抗糖尿病潜力和油性石楠(卢尔) 果实的超微量色谱指纹图谱. *生药学杂志*, 2019, 11(3): 469-474. <https://doi.org/10.5530/pj.2019.11.74>
- [12] HANIFAH L. 黑桑叶提取物 (油菜花) 对体外增殖和细胞凋亡率 (细胞核) 的影响. *期刊生物群*, 2020, 6 (1): 19-25. <https://doi.org/10.19109/biota.v6i1.4372>
- [13] DEWI Y. K. 使用多级提取方法对黑桑叶树皮提取物 [油菜花(卢尔) 果] 进行抗氧化活性测试. *旋转. 化学期刊和彭迪次基米亚*, 2021, 3(1): 22-31. <https://doi.org/10.20414/spin.v3i1.3020>
- [14] YANG WANG T., LI Q., 和 SHUN BI K. 药用植物中的生物活性黄酮类化合物: 结构、活性和生物命运. *亚洲药理学杂志*, 2018, 13(1): 12-23. <https://doi.org/10.1016/j.ajps.2017.08.004>
- [15] CARITÀ A. C., FONSECA-SANTOS B., SHULTZ J. D., MICHNIAK-KOHN B., CHORILLI M. 和 LEONARDI G. R. 维生素 C: 一种化合物, 多种用途. 交付、效率和稳定性方面的进步. *纳米医学: 纳米技术、生物学和医学*, 2020, 24: 102117. <https://doi.org/10.1016/j.nano.2019.102117>
- [16] IFEANYI O. E. 自由基和抗氧化剂综述. *国际医学科学期刊*, 2018, 4(2): 123-133. <http://dx.doi.org/10.22192/ijcrms.2018.04.02.019>

- [17] ALKADI H. 自主基和抗氧化剂途径。糖尿病。药物观点，2018，20(1)：16-26。
<https://doi.org/10.2174/1871526518666180628124323>
- [18] MCIVER L. A. 和 TRIPP J. 继续教育活动。统计珍珠出版，国家生物技术信息中心，2021。
<https://www.ncbi.nlm.nih.gov/books/NBK493214>
- [19] VIEIRA R.，SOUTO S. B.，SÁNCHEZ-LÓPEZ E.，MACHADO A. L.，SEVERINO P.，JOSE S.，SANTINI A.，FORTUNA A.，GARCÍA M. L.，SILVA A. M.，和 SOUTO E. B. 糖尿病药物 2 型糖尿病和代谢综合征——经共价化合物综述：第一部分。制药，2019，12(4)：152。
<https://doi.org/10.3390/ph12040152>
- [20] GOSWAMI S.，& SINGH R. P. 阿育吠陀：油桐花(劳尔)。美药：一种具有巨大生物活性的传统植物的植物化学和药理学讨论。世界药物研究杂志，2017，6(10)：295-309。
<https://doi.org/10.20959/wjpr201710-9370>
- [21] JOSE S.，& SINGH R. P. 杜松花和油菜籽的水性和甲醇提取物的抗氧化能力比较分析。巴尔尼德研究杂志，2018，9(3)：221-227。
<https://doi.org/10.12680/balneo.2018.185>
- [22] SHANIN S. A. H.，NABAT Z. M.，SALIH M. A. 和 AL KHAFAH B. J. 使用扩展二进制布谷鸟搜索和克-最近邻分类器诊断糖尿病。西贡交通大学学报，2020，55(2)。
<https://doi.org/10.35741/issn.0258-2724.55.2.36>
- [23] ŞÖHRETOĞLU D.，& SARI S. 黄酮类化合物作为 α -葡萄糖苷酶抑制剂：机械方法与酶动力学和分子建模相结合。植物化学评论，2020，19(5)：1081-1092。
<https://doi.org/10.1007/s11101-019-09610-6>

JURNAL SCOPUS 2

ORIGINALITY REPORT

23%
SIMILARITY INDEX

18%
INTERNET SOURCES

17%
PUBLICATIONS

4%
STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | www.id-press.eu
Internet Source | 3% |
| 2 | Yeni Susanti, Merryana Adriani, Annis Catur Adi. "Effect of Single Clove Garlic Extract (Allium Sativum Linn) on Blood Sugar Levels, Malondialdehyde, Insulin Levels and Insulin Resistance (Experiments in Rats (Rattus Novergicus) Induced by Streptozotocin", STRADA Jurnal Ilmiah Kesehatan, 2020
Publication | 2% |
| 3 | m.scirp.org
Internet Source | 2% |
| 4 | www.riset.unisma.ac.id
Internet Source | 1% |
| 5 | www.gssrr.org
Internet Source | 1% |
| 6 | R R Setiadi, A F M Z Zein, D Nauphar. "Antihyperglycemic effectiveness comparison of ethanol extract of soursop leaf (L.) againsts acarbose in streptozotocin-induced diabetic | 1% |

white rats ", Journal of Physics: Conference
Series, 2019

Publication

7	jurnal.radenfatah.ac.id Internet Source	1 %
---	--	-----

8	www.e3s-conferences.org Internet Source	1 %
---	--	-----

9	Submitted to Universitas Diponegoro Student Paper	1 %
---	--	-----

10	Zaldy Rusli, Bina Lohita Sari, Novi Fajar Utami, S Sabila. "Optimization Of Microwave-Assisted Extraction Of Flavonoids From Binahong (Anredera cordifolia) Leaves Using Respon Surface Methodology", Jurnal Fitofarmaka Indonesia, 2020 Publication	1 %
----	---	-----

11	Purnama Ningsih, Sitti Rahmawati, Baharuddin Hamzah, Tri Santoso, Nurbaya Nurbaya, Muhammad Fakhrul Hardani, Ririen Hardani. "Histology of hematoxylin–eosin and immunohistochemical diabetes rat pancreas after giving combination of moringa leaves (Moringa oleifera) and clove flower (Syzygium aromaticum) extracts", Open Access Macedonian Journal of Medical Sciences, 2021 Publication	1 %
----	--	-----

12	leju.eu Internet Source	
----	----------------------------	--

1 %

13

ejournal.unsri.ac.id

Internet Source

<1 %

14

scitepress.org

Internet Source

<1 %

15

www.dovepress.com

Internet Source

<1 %

16

Wahyudin ., Muh Nasrum Massi, Rosdiana Natzir, Gemini Alam, Agus Salim Bukhari. "Effect of Sukun Leaf Extract [Artocarpus altilis (Park.) Fosberg] on Insulin Resistance in Obese Rats (Rattus norvegicus): A Study of Free Fatty Acid (FFA) Levels", Pakistan Journal of Nutrition, 2017

Publication

<1 %

17

Afsaneh Pashapoor, Seeyamak Mashhadyrafie, Pejman Mortazavi. "Ameliorative effect of Myristica fragrans (nutmeg) extract on oxidative status and histology of pancreas in alloxan induced diabetic rats", Folia Morphologica, 2015

Publication

<1 %

18

Submitted to University of Malaya

Student Paper

<1 %

19 Montanya, E.. "Islet- and stem-cell-based tissue engineering in diabetes", Current Opinion in Biotechnology, 200410 <1 %
Publication

20 zh-tw.top10supps.com <1 %
Internet Source

21 "Abstracts of the EASD, Stockholm 2010", Diabetologia, 2010 <1 %
Publication

22 Submitted to School of Business and Management ITB <1 %
Student Paper

23 Yuliet G. Latuhihin, Theopilus Watuguly, Pieter Kakisina, Indranila Kustarini Samsuria. "Potential of Binahong (*Anredera cordifolia* [Tenore] Steen) in Reducing TNF- α Expression on Regeneration of Pancreas β Cells on White Rats (*Rattus norvegicus*) Diabetes Mellitus Models", Journal of Biosciences and Medicines, 2020 <1 %
Publication

24 doaj.org <1 %
Internet Source

25 ejournal.umm.ac.id <1 %
Internet Source

26 www.buy123.com.tw <1 %
Internet Source

27 www.x-mol.com <1 %
Internet Source

28 "23rd Annual Meeting of the European Association for the Study of Diabetes Leipzig, German Democratic Republic, 15-19 September 1987", Diabetologia, 1987
Publication

29 Ali Muhammed Moula Ali, Maria Gullo, Amit Kumar Rai, Sri Charan Bindu Bavisetty. "Bioconservation of iron and enhancement of antioxidant and antibacterial properties of chicken gizzard protein hydrolysate fermented by 8042 ", Journal of the Science of Food and Agriculture, 2020
Publication

30 ajtr.org <1 %
Internet Source

31 journal.uinmataram.ac.id <1 %
Internet Source

32 pdf.aminer.org <1 %
Internet Source

33 plantarchives.org <1 %
Internet Source

wrhrc.tbzmed.ac.ir

34

Internet Source

<1 %

35

baadalsg.inflibnet.ac.in

Internet Source

<1 %

36

bianke.cnki.net

Internet Source

<1 %

37

caod.oriprobe.com

Internet Source

<1 %

38

czkj.ci.cqvip.com

Internet Source

<1 %

39

Ganji Purnachandra Nagaraju, Syed F Zafar, Bassel F El-Rayes. "Pleiotropic effects of genistein in metabolic, inflammatory, and malignant diseases", Nutrition Reviews, 2013

Publication

<1 %

40

Suo-Lin Fu, Hui-Ming Zhu. "Surgical treatment of morbid obesity and type 2 diabetes mellitus: current situation and future prospects", World Chinese Journal of Digestology, 2010

Publication

<1 %

41

dspace.xmu.edu.cn

Internet Source

<1 %

42

eprints.whiterose.ac.uk

Internet Source

<1 %

43	google.com.na Internet Source	<1 %
44	journals.plos.org Internet Source	<1 %
45	theodbo-korea.blogspot.com Internet Source	<1 %
46	www.24drs.com Internet Source	<1 %
47	www.iosrjournals.org Internet Source	<1 %
48	zl50.com Internet Source	<1 %
49	Elly Tugiyanti, Emmy Susanti, Ibnu Hari Sulis. "Effect of Tea Dregs Form and Different Fermentation Process on the Nutrient, Tannin, Saponin, flavonoid content and Antioxidant Activity", Pakistan Journal of Nutrition, 2019 Publication	<1 %
50	Hussam M. Gharib, Mohammad Y. Abajy, Abdulnaser Omaren. "Investigating the effect of some fluoroquinolones on C-reactive protein levels and ACh-Induced blood pressure reduction deviations after aging of diabetes in STZ-Induced diabetic wistar rats", Heliyon, 2020	<1 %

51 Piao Wang, Yi Liu, Tong Zhang, Cheng Yin, Seok Yong Kang, Su Jin Kim, Yong-Ki Park, Hyo Won Jung. " Effects of Root Extract of in Mice with High-Fat-Diet/Streptozotocin-Induced Diabetes and C2C12 Myoblast Differentiation ", ACS Omega, 2021

Publication

<1 %

52 Submitted to Universitas Hasanuddin

Student Paper

<1 %

53 cofacts.g0v.tw

Internet Source

<1 %

54 dict.cnki.net

Internet Source

<1 %

55 ejournal.uin-malang.ac.id

Internet Source

<1 %

56 genewei.com

Internet Source

<1 %

57 juniperpublishers.com

Internet Source

<1 %

58 rasayanjournal.co.in

Internet Source

<1 %

59 www.semanticscholar.org

Internet Source

<1 %

60

B. Kojić-Prodić, Ž. Ružić-Toroš, D. Grdenić, L. Golič. " The crystal structure of dioxobis(1,3-diphenylpropanedionato)molybdenum(VI), (C₁₂H₁₀O₄)MoO₄ ", Acta Crystallographica Section B Structural Crystallography and Crystal Chemistry, 1974

Publication

<1 %

61

E Peniati, E Setiadi, R Susanti, R S Iswari. " Anti-hyperglycemic effect of peel extract on blood sugar level of alloxan-induced Wistar rats ", Journal of Physics: Conference Series, 2018

Publication

<1 %

62

Geon-Woo Kim, Yun-Hee Choi, Byung-Loc Kim, Yonguk Kim, Rack-Seon Seong, Min-Hee Han, Gyeong-Ae Kim, Min-Ju Choi, Yong-Gi Jeong. "Determination of Anti-oxidative and Whitening Effects of Complex Extracts Obtained from Sprout Panax ginseng C.A. Meyer and Cassia nomame (Sieb.) Honda on Skin", Asian Journal of Beauty and Cosmetology, 2018

Publication

<1 %

63

Juan Zhang, Jia-Zhao Fu, Shi-Fu Hong, Hong Jiang, Zhong-Quan Qi, Zhao-Sui Huang, Jun-Jie Xia. "Toxicity of rapamycin and its derivatives to pancreatic islets", World Chinese Journal of Digestology, 2016

<1 %

64

R D Andriani, P P Rahayu, F Chairunnisa, S N Wulan, J M Maligan. "Fortification of skim milk with whey protein xanthone and its effect on antihyperglycemic activities in animal model", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

65

K Tsukiyama. "Gastric inhibitory polypeptide is the major insulinotropic factor in K(ATP) null mice", European Journal of Endocrinology, 09/01/2004

Publication

<1 %

66

Letícia Aparecida Ferreira de Abreu, Renato Paiva, Judith Georgette Alcalde Mosqueira, Michele Valquíria dos Reis et al. "Antioxidant activity and physico-chemical analysis of Campomanesia rufa (O.Berg) Nied. fruits", Ciência e Agrotecnologia, 2020

Publication

<1 %

67

Zong-Ting Gu, Zong-Ze Li, Cheng-feng Wang. "Research advances of intracellular mechanisms underlying gemcitabine resistance in pancreatic cancer", World Chinese Journal of Digestology, 2020

Publication

<1 %

68

www.taiwanpropolis.com
Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On